

Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

DYER. The third supplement appears under the direction of D. Prain, the new director of the Kew Gardens. The supplement is a record of the remarkable activity in taxonomy during the five years covered, and critical judgment in reference to this great volume of work is becoming increasingly difficult. In fact, the list is a record of publication rather than an expression of opinion. For example, 476 acknowledged species of Crataegus are recorded for the five years, and Rubus is not far behind. To review such a work is impossible. It is only necessary to announce its appearance.—J. M. C.

North American Flora.—The third part of volume 22 has just been issued. It contains Grossulariaceae by F. V. Coville and N. L. Britton, 43 species (2 new) being referred to Ribes and 40 (4 new) to Grossularia; Platanaceae (6 spp.) by H. A. Gleason; Crossosomataceae (4 spp., 1 new) by J. H. Small; Connaraceae (3 genera, 9 spp.) by N. L. Britton; Calycanthaceae (4 spp.) by C. L. Potter; and the beginning of Rosaceae by P. A. Rydberg, the key recognizing 18 tribes, 6 of which are completed and the seventh (Potentilleae) begun. Among the 19 genera of Rosaceae presented, Horkeliella (3 spp.) is new; and 29 new species are described, being distributed among Opulaster (4), Spiraea (5), Petrophytum, Aruncus (3), Chamaebatiaria, Lindleyella, Vauquelinia, Sericotheca (6), Horkelia (6), and Ivesia.—J. M. C.

Marine algae of Sweden.—KYLIN⁷ has published a monograph of the algal flora of the west coast of Sweden. The species of the four following groups are first presented: Chlorophyceae (12 fams., 26 gen., 71 spp.), Fucoideae (16 fams., 51 gen., 105 spp.), Bangiales (5 gen., 11 spp.), Florideae (16 fams., 55 gen., 107 spp.). One new genus (*Acrothrix*) of Fucoideae is described. The second part of the contribution (80 pp.) presents the ecological factors and analyses the geographical distribution. At the conclusion of the discussion, the 105 species of Fucoideae and the 118 species of Florideae (incl. Bangiales) are distributed into arctic, subarctic, boreal-arctic, cold-boreal, and warm-boreal groups. The paper concludes with "biological" notes, a full bibliography, and an adequate index.—J. M. C.

Das Pflanzenreich.⁸—Part 33 contains the 9 genera of Aloineae (Liliaceae) by A. Berger, *Chortolirion* being a new genus with 4 species. Altogether, 370 species are presented, many of them with numerous cultivated forms and entering into hybrids. The large genera are Aloe (178 spp., 14 new), Kniphofia (67 spp., 2 new), Haworthia (60 spp.), and Gasteria (43 spp.).

Part 34 contains the Sarraceniaceae by J. M. MACFARLANE, who gives an interesting account (in English) of the structure of the vegetative organs and the

⁷ KYLIN, HARALD, Studien über die Algenflora der schwedischen Westküste. pp. 288. pls. 7. Upsala. 1907. (Inaugural dissertation.)

⁸ ENGLER, A., Das Pflanzenreich. Heft 33, Liliaceae-Asphodeloideae-Aloineae von A. Berger. pp. 347. figs. 141 (817). M17.60. Heft 34, Sarraceniaceae von J. M. MACFARLANE. pp. 39. figs. 10 (43). M2.40. Leipzig: Wilhemn Engelmann. 1908.